

On the integrability of spherical gravitational waves in vacuum

Ioannis Bakas*

*Theory Division, Department of Physics, CERN
CH-1211 Geneva 23, Switzerland*

`ioannis.bakas@cern.ch`

Abstract

The general class of Robinson-Trautman metrics that describe gravitational radiation in the exterior of bounded sources in four space-time dimensions is shown to admit zero curvature formulation in terms of appropriately chosen two-dimensional gauge connections. The result, which is valid for either type II or III metrics, implies that the gravitational analogue of the Lienard-Wiechert fields of Maxwell equations form a new integrable sector of Einstein equations for any value of the cosmological constant. The method of investigation is similar to that used for integrating the Ricci flow in two dimensions. The zero modes of the gauge symmetry (factored by the center) generate Kac's K_2 simple Lie algebra with infinite growth.

*On sabbatical leave from Department of Physics, University of Patras, GR-26500 Patras, Greece;
e-mail: bakas@ajax.physics.upatras.gr

The theory of gravitational waves is an old subject. The existence of general asymptotically flat radiative space-times and the explicit construction of various classes of solutions have received considerable attention over the years. They have several applications that serve as testing bed for the physical properties of gravitation. Spherical gravitational waves are quite interesting in this respect, as they are thought to represent an isolated gravitationally radiating system. Yet, there has been no systematic way to understand the algebraic structure of the non-linear equations that govern their propagation in vacuum, apart from global considerations and the behavior of solutions in the asymptotic future. A few explicit solutions are also known to this day. It is the purpose of the present work to provide a new formulation of the problem by casting the corresponding sector of Einstein equations into zero curvature form, thus making a decisive step toward their integration. The mathematical framework that achieves this purpose points to the relevance of a novel class of infinite dimensional Lie algebras that might be of more general value in dynamical gravitational problems.

Recall that the most general gravitational vacuum solution in four space-time dimensions which admits a geodesic, shear-free, twist-free (but diverging) null congruence is described by the class of Robinson-Trautman metrics with line element, [1, 2],

$$ds^2 = 2r^2 e^\Phi dz d\bar{z} - 2dt dr - H dt^2 , \quad (1)$$

where $H(z, \bar{z}; t)$ has the special form

$$H = r \partial_t \Phi - \Delta \Phi - \frac{2m(t)}{r} - \frac{\Lambda r^2}{3} . \quad (2)$$

The affine parameter r varies along the rays of the repeated null eigenvector and t is a retarded time coordinate. Closed surfaces of constant r and t represent distorted two-dimensional spheres Σ with metric coefficient given by $\Phi(z, \bar{z}; t)$ in the system of conformally flat coordinates $(z, \bar{z})$. The parameter $m(t)$, which in some cases represents the physical mass of the system, can be set equal to a constant when differs from zero; in the sequel m is taken positive and set equal to $1/3$ by appropriate relabeling of the null hyper-surfaces in the Robinson-Trautman class of metrics. The special case $m = 0$ will be considered separately at the end. Finally, Λ is the value of the cosmological constant in space-time, which can be positive, negative or zero depending on the physical circumstances.

There is a close analogy between the Lienard-Wiechert fields of Maxwell equations and the Robinson-Trautman metrics of Einstein equations for they both admit a principal null vector field which is geodesic, shear and twist free with non-vanishing divergence, thus giving a more intuitive meaning to the general ansatz for the metrics above, [3]. In the gravitational case, the field equations imply that all vacuum solutions in this class with cosmological constant Λ satisfy the following parabolic fourth order differential equation for the unknown function Φ ,

$$\partial_t \Phi = -\Delta \Delta \Phi , \quad (3)$$

where $\Delta = \exp(-\Phi) \partial \bar{\partial}$ denotes the Laplace-Beltrami operator on the distorted two-dimensional spheres Σ . Note that this non-linear equation, which is named after Robinson

and Trautman, is not sensitive to the value of Λ . Its time dependent solutions can be employed for the propagation of spherical gravitational waves in vacuum, although the strict sense of the word spherical symmetry will automatically imply that the metric is static. Thus, it is more appropriate to think of such metrics as describing gravitational radiation outside some bounded region.

Its fixed point solutions include the static Schwarzschild metric with mass m (written in Eddington-Filkenstein frame) to which infinitely many trajectories tend in the asymptotic future. Approximate solutions were first obtained by perturbing the Schwarzschild metric so that the geometry of the two-dimensional slices Σ represents distorted (e.g., axially symmetric) configurations in multi-pole expansion. It follows, in the linearized approximation to equation (3), that small perturbations associated to spherical harmonics with angular momentum $l \geq 2$ decay exponentially fast by inducing quadrupole and higher order pole radiation, and the configuration settles back to the round sphere as $t \rightarrow \infty$, [4]. Subsequently, it was found that all dynamical solutions exhibit Lyapunov stability using the quadratic curvature functional

$$L(t) = \int_{\Sigma} R^2 d\sigma , \quad (4)$$

where R is the Ricci scalar curvature of Σ and $d\sigma$ its volume element, [5]. $L(t)$ decreases monotonically with time and the critical point, where it assumes its extremal value, is the round sphere corresponding to the Schwarzschild solution. Finally, the (semi)-global existence and convergence of solutions with smooth initial data has been established in all generality, following similar arguments to the long time existence and convergence of solutions to the normalized Ricci flow, thus establishing the universal limiting character of all asymptotically flat Robinson-Trautman metrics, [6]; see also references [7] for related earlier work. Generalizations to space-times with non-vanishing cosmological constant have also been considered in the literature, [8].

The Robinson-Trautman equation also arose independently in mathematics while considering the notion of extremal metrics on compact Kähler spaces K , [9]. Using Kähler coordinates $(z_a, \bar{z}_a)$, the following deformation equation was introduced for the metric $g_{a\bar{b}}$ with Ricci scalar curvature R ,

$$\partial_t g_{a\bar{b}} = \partial_a \bar{\partial}_b R . \quad (5)$$

The quadratic curvature functional (4), which is in general defined over the Kähler manifold K , decreases monotonically under the flow and its minimum defines extremal metric on it. When certain obstructions vanish, [10], the extremal metric is canonical in that its curvature is constant. The equation (5) is also known in the literature as Calabi flow and for one-dimensional complex Kähler spaces, in particular, it coincides with the main equation (3), as first noted in reference [11]. In this context, the convergence of the Robinson-Trautman metrics to the Schwarzschild solution in the asymptotic future implies that the extremal metric on a two-dimensional sphere is the round one.

Next, we sketch the algebraic framework that casts the Robinson-Trautman equation into zero curvature form. Let us introduce an infinite dimensional algebra whose local

part is spanned by six different elements $H_i(t)$, $X_i^\pm(t)$ with the index i taking the values 0 and 1. These generators depend on a continuous real variable t and assumed to obey the basic system of commutation relations

$$\begin{aligned} [H_0(t), X_0^\pm(t')] &= \pm\delta(t-t')X_0^\pm(t') , & [X_1^\pm(t), X_0^\mp(t')] &= \delta(t-t')H_0(t') , \\ [H_0(t), X_1^\pm(t')] &= \pm\delta(t-t')X_1^\pm(t') , & [X_1^+(t), X_1^-(t')] &= c\delta(t-t')H_1(t') , \\ [H_1(t), X_1^\pm(t')] &= -\partial_t\delta(t-t')X_0^\pm(t') , \end{aligned} \quad (6)$$

where c is an arbitrary real constant; its physical meaning will become clear later. The remaining set of basic commutation relations are taken to be trivial, that is impose

$$\begin{aligned} [H_1(t), X_0^\pm(t')] &= 0 , & [X_0^+(t), X_0^-(t')] &= 0 , \\ [H_i(t), H_j(t')] &= 0 . \end{aligned} \quad (7)$$

It can be verified that the commutation relations above are consistent with the Jacobi identities for all values of c .

The algebra can be enlarged further by taking successive commutators among its basic generators $X_i^\pm$, such as $[\dots[X_i^\pm, X_j^\pm], \dots, X_k^\pm]$, that give rise to new elements, as in the simpler case of Kac-Moody algebras where $X_i^\pm$ act as creation and annihilation operators in the Verma module representations; in that context H_i are the Cartan generators of the commutative subalgebra. The structure of the higher commutation relations can be in principle established working order by order and requiring consistency with the Jacobi identities. However, there are compelling reasons to believe that the complete characterization of all elements of the algebra at hand, together with their commutations relations, is a very difficult task. As will be seen later, its zero modes (factored by the center) generate Kac's K_2 algebra, which is the most elementary example of infinite dimensional simple Lie algebras with infinite growth. Also, a simpler version of this algebra will be encountered later in a different - yet related problem - and it grows exponentially fast beyond the local part. Despite these limitations, which are also common to hyperbolic Kac-Moody algebras, the zero curvature formulation of the Robinson-Trautman equation can be established by choosing gauge connections with values only in the local part.

An alternative description of the same algebraic structure is obtained by smearing all generators by appropriate smooth functions of compact support, $\varphi(t)$, and integrating over the entire range of t ,

$$A(\varphi) = \int \varphi(t) A(t) dt , \quad (8)$$

as commonly used in the theory of distributions and in the current algebras of quantum field theory. Then, the above system of commutation relation takes the form

$$\begin{aligned} [H_0(\varphi), X_0^\pm(\psi)] &= \pm X_0^\pm(\varphi\psi) , & [X_1^\pm(\varphi), X_0^\mp(\psi)] &= H_0(\varphi\psi) , \\ [H_0(\varphi), X_1^\pm(\psi)] &= \pm X_1^\pm(\varphi\psi) , & [X_1^+(\varphi), X_1^-(\psi)] &= c H_1(\varphi\psi) , \\ [H_1(\varphi), X_1^\pm(\psi)] &= X_0^\pm(\varphi'\psi) , \end{aligned} \quad (9)$$

and

$$\begin{aligned} [H_1(\varphi), X_0^\pm(\psi)] &= 0, & [X_0^+(\varphi), X_0^-(\psi)] &= 0, \\ [H_i(\varphi), H_j(\psi)] &= 0. \end{aligned} \quad (10)$$

The prime here denotes derivative with respect to the continuous index t .

It is more convenient to use the latter form and introduce gauge connections on the two-dimensional space Σ taking values in this infinite dimensional algebra. Postulating the zero curvature condition

$$F_{z\bar{z}} = [\partial + A_+(z, \bar{z}; t), \bar{\partial} + A_-(z, \bar{z}; t)] = 0, \quad (11)$$

where the two components of the gauge connection $A_\pm$ are taken to depend on four functions f, g, Φ and Ψ of $z, \bar{z}$ and t in the following way,

$$\begin{aligned} A_+ &= H_0(f) + H_1(g) + X_1^+(1), \\ A_- &= X_0^-(\Psi e^\Phi) + X_1^-(e^\Phi), \end{aligned} \quad (12)$$

we arrive at a consistent system of coupled differential equations for the unknown functions. The computation of $F_{z\bar{z}}$ gives rise to various terms taking values in linear span of the generators H_0, H_1, X_1^- and X_0^- alone. These are required to vanish separately by the zero curvature condition (11) and they lead, respectively, to the four equations

$$\begin{aligned} \bar{\partial}f &= \Psi e^\Phi, & \bar{\partial}g &= e^\Phi, & \partial\Phi &= f, \\ \partial(\Psi e^\Phi) - f\Psi e^\Phi + (\partial_t g)e^\Phi &= 0 \end{aligned} \quad (13)$$

provided that the constant c in the commutation relations is chosen to be 1.

Eliminating $f = \partial\Phi$ from the first equation, it follows that

$$\partial\bar{\partial}\Phi = \Psi e^\Phi. \quad (14)$$

Also, eliminating f from the very last equation, it follows that $\partial_t g = -\partial\Psi$, which upon substitution into the t -derivative of second equation yields

$$\partial_t e^\Phi = -\partial\bar{\partial}\Psi. \quad (15)$$

Note that according to equation (14), the variable Ψ is minus the Ricci scalar curvature of the two-dimensional line element on Σ ,

$$ds^2 = 2e^\Phi dzd\bar{z}, \quad (16)$$

which has spherical topology and sits inside the Robinson-Trautman metric. When Ψ is eliminated from the combined system (14) and (15), one readily obtains $\partial_t\Phi = -\Delta\Phi$, which is the same as (3). Thus, the Robinson-Trautman equation arises as zero curvature condition $F_{z\bar{z}} = 0$ on the two-dimensional space of constant (r, t) slices. The infinite

dimensional algebra where the gauge connections assume their values is quite novel and has not been considered in the physics or mathematics literature before, to the best of our knowledge. It certainly deserves further study in order to be able to construct explicitly conserved integrals of the equation and present its general solution by Bäcklund transformations.

The situation is reminiscent, though is more complicated now, to the algebraic description of the Ricci flow in two dimensions using conformally flat (Kähler) coordinates. In that case, the evolution equation $\partial_t g_{z\bar{z}} = -R_{z\bar{z}}$ for the t -dependent class of metrics (16) reads $\partial_t \Phi = \Delta \Phi$. As shown in [12, 13], the Ricci flow in two dimensions admits a zero curvature formulation, as in equation (11) above, choosing two-dimensional gauge connections of the form

$$A_+ = H(\Psi) + X^+(1) , \quad A_- = X^-(e^\Phi) , \quad (17)$$

where $\partial_t \Psi = -\partial \Phi$. It was further assumed that the elements $H, X^\pm$ generate the local part of an infinite dimensional Lie algebra with basic commutation relation given (in smeared form) by

$$\begin{aligned} [X^+(\varphi), X^-(\psi)] &= H(\varphi\psi) , & [H(\varphi), H(\psi)] &= 0 , \\ [H(\varphi), X^\pm(\psi)] &= \pm X^\pm(\varphi'\psi) . \end{aligned} \quad (18)$$

They provide the local part of a novel infinite dimensional algebra with anti-symmetric Cartan kernel $K(t, t') = \partial_t \delta(t - t')$, due to the parabolic nature of the Ricci flow in time, whose completion by successive commutation of X^+ 's and X^- 's yields a graded algebra with exponentially fast growth, [14].

The Robinson-Trautman equation or Calabi flow, as it is also known in the mathematics literature, is a parabolic fourth order differential equation that is formally related to the second order Ricci flow by simply taking the square root of the t -derivative operator. It can be easily seen how this procedure works and changes the order of the equations accordingly. Consider, in particular, the abstract super-evolution Ricci type flow

$$\partial \bar{\partial} \mathcal{F} = \mathcal{D} \exp \mathcal{F} , \quad (19)$$

where $\mathcal{F}(z, \bar{z}; t, \theta) = \Phi(z, \bar{z}; t) + \theta \Psi(z, \bar{z}; t)$ is a mixed superfield with bosonic components Φ and Ψ and $\mathcal{D} = \partial/\partial\theta - \theta\partial/\partial t$ is the superderivative in $(1, 1)$ superspace with coordinates (t, θ) , so that $\theta^2 = 0$ and $\mathcal{D}^2 = -\partial/\partial t$. Expanding $\exp \mathcal{F} = (1 + \theta \Psi) \exp \Phi$ and comparing the even and odd parts of the super-flow (19), one immediately arrives at the system (14) and (15) that amount to the Robinson-Trautman equation for the variable Φ . Note on the side that the procedure of taking the square root of the time derivative operator might be of more general value relating other pairs of second and fourth order parabolic equations.

According to this observation, it is natural to expect that there is an infinite dimensional superalgebra at work, which generalizes the commutation relations (18) and

accounts for the (seemingly ad hoc) choice of the bosonic algebra (9) and (10) used in the theory of spherical gravitational waves. By the same token, it is also natural to expect that the general solution can be obtained in closed form using group theoretical methods similar to those employed for the Ricci flow, [12, 13]. It should be emphasized in both cases that the parabolic nature of the equations, which tend to dissipate curvature perturbations from the canonical form of the metric, is not in contradiction with their integrability. The latter refers to their zero curvature description in two dimensional spaces with local coordinates $(z, \bar{z})$, whereas the deformation variable t is treated differently by accommodating it into the structure of the Lie algebra where the gauge connections take values. The reciprocity of our description is common to other examples of non-linear differential equations. Note, however, that the underlying algebras do not necessarily possess a non-zero invariant bilinear form, in contrast to the case of Kac-Moody algebras with symmetrizable Cartan matrices. Thus, the exact mathematical meaning of integrability and the construction of conserved integrals in two dimensions present some challenging questions. Further details on these issues will be given elsewhere, [15].

We turn now to the special case of Robinson-Trautman metrics with mass parameter $m = 0$ for which the main equation (3) simplifies to

$$\Delta\Delta\Phi = 0 . \quad (20)$$

Equivalently, it can be stated as $\Delta\Phi(z, \bar{z}; t) = \chi(z, t) + \bar{\chi}(\bar{z}; t)$, where χ and $\bar{\chi}$ are holomorphic and anti-holomorphic functions, respectively, that are complex conjugate to each other. Note here that t appears as spectator with no apparent role in the dynamics. This class of solutions is naturally related to Nirenberg's mathematical problem for finding metrics of prescribed curvature on S^2 , since $R[S^2] = -\Delta\Phi$; see, for instance, [16], for a recent approach based on flows. The zero curvature formulation of equation (20) is slightly different as it requires the choice of gauge connections (12) taking values in the infinite dimensional algebra (9), (10) with parameter $c = 0$. This, in turn, implies that the resulting system is the same as in equation (13) given above, apart from the component $\bar{\partial}g = \exp\Phi$ that is now replaced by $\bar{\partial}g = 0$. Eliminating f and g , as before, it follows

$$\partial\bar{\partial}\Phi = \Psi e^\Phi , \quad \partial\bar{\partial}\Psi = 0 \quad (21)$$

that yield equation (20) as required. Finally, note that the one-parameter family of holomorphic functions g and χ are simply related to each other by $\partial\chi = -\partial_t g$.

In conclusion, it is instructive to compare our results with other works on the integration of the Robinson-Trautman equation by prolongation methods. Type II solutions (in terms of Petrov classification), which are the generic examples when the mass parameter $m \neq 0$, have resisted all previous attempts with the exception of some partial (yet inconclusive) results described in reference [17]. On the other hand, type III solutions, which have $m = 0$, have been more tractable by such methods. The corresponding metrics were shown to exhibit a rich prolongation structure given by the simple infinite dimensional algebra K_2 (in Kac's notation), [18]. This result fits nicely into the algebra (9), (10)

with $c = 0$, since the zero modes of the currents $H_i(t)$, $X_i^\pm(t)$ form the local part of K_2 algebra. Introducing the change of basis

$$\begin{aligned} e_0 &= \frac{1}{\sqrt{2}}(X_0^+ + X_1^+) , & e_1 &= \frac{1}{\sqrt{2}}(-X_0^+ + X_1^+) , \\ f_0 &= \frac{1}{\sqrt{2}}(X_0^- - X_1^-) , & f_1 &= \frac{1}{\sqrt{2}}(X_0^- + X_1^-) , \end{aligned} \quad (22)$$

it easily follows that the zero modes of the currents, without t -dependence, satisfy the basic system of commutation relations

$$[e_i, f_j] = \delta_{ij}h , \quad [h, e_i] = e_i , \quad [h, f_i] = -f_i \quad (23)$$

with h given by the zero mode of H_0 . These are precisely the defining relations for the local part of the algebra K_2 , [19], whereas the zero mode of the generator H_1 commutes with the rest and decouples.

For type II metrics with mass parameter m , the commutation relations (9), (10) have $c = 3m$, in which case the zero mode of the current H_1 generates the center of the algebra with coefficient (central charge) c . Thus, K_2 also describes the algebra of the zero modes of the currents in type II situations provided that the one-dimensional center is factored out. In either case, the infinite dimensional algebra we introduced in this paper is much larger, as it includes an infinite number of Fourier modes, and (apparently) provides the minimal choice for establishing the zero curvature formulation of the Robinson-Trautman equation for spherical gravitational waves in vacuum for all values of the mass parameter and the cosmological constant.

It will be interesting to see how the results may generalize to Einstein-Maxwell theory. Particular interest also present the generalizations in the presence of additional massless fields that arise in string theory, [20], and their possible implications for the gravity/gauge correspondence beyond the plane wave limit.

Acknowledgments

This work was supported in part by the European Research and Training Network “Constituents, Fundamental Forces and Symmetries of the Universe” under contract number MRTN-CT-2004-005104 and the INTAS program “Strings, Branes and Higher Spin Fields” under contract number 03-51-6346. I thank Ecole Polytechnique for hospitality in the initial stages of the present work and CNRS for financial support. I also thank the Theory Division at CERN for hospitality and financial support during my sabbatical leave in the academic year 2004-05, where the main body of the present work was carried out in excellent and stimulating environment.

References

- [1] I. Robinson and A. Trautman, “Spherical gravitational waves”, *Phys. Rev. Lett.* **4** (1960) 431; “Some spherical gravitational waves in general relativity”, *Proc. Roy. Soc. A265* (1962) 463.
- [2] D. Kramer, H. Stephani, E. Herlt and M. MacCallum, *Exact Solutions of Einstein’s Field Equations*, Cambridge University Press, 1980.
- [3] E.T. Newman, “Lienard-Wiechert fields and general relativity”, *J. Math. Phys.* **15** (1974) 44.
- [4] J. Foster and E.T. Newman, “Note on the Robinson-Trautman solutions”, *J. Math. Phys.* **8** (1967) 189.
- [5] B. Lukacs, Z. Perjes, J. Porter and A. Sebestyen, “Lyapunov functional approach to radiative metrics”, *Gen. Relat. Grav.* **16** (1984) 691.
- [6] P. Chrusciel, “Semi-global existence and convergence of solutions of the Robinson-Trautman (2-dimensional Calabi) equation”, *Commun. Math. Phys.* **137** (1991) 289; “On the global structure of Robinson-Trautman space-times”, *Proc. Roy. Soc. A436* (1992) 299.
- [7] B. Schmidt, “Existence of solutions of the Robinson-Trautman equation and spatial infinity”, *Gen. Relat. Grav.* **20** (1988) 65; A. Rendall, “Existence and asymptotic properties of global solutions of the Robinson-Trautman equation”, *Class. Quant. Grav.* **5** (1988) 1339; D. Singleton, “On global existence and convergence of vacuum Robinson-Trautman solutions”, *Class. Quant. Grav.* **7** (1990) 1333.
- [8] J. Bicak and J. Podolsky, “Global structure of Robinson-Trautman radiative space-times with a cosmological constant”, *Phys. Rev.* **D55** (1997) 1985.
- [9] E. Calabi, “Extremal Kähler metrics”, in *Seminar on Differential Geometry*, ed. S.-T. Yau, *Annals of Mathematics Studies*, vol. 102, Princeton University Press, 1982.
- [10] A. Futaki, “An obstruction to the existence of Einstein Kähler metrics”, *Invent. Math.* **73** (1983) 437.
- [11] K.P. Tod, “Analogues of the past horizon in the Robinson-Trautman metrics”, *Class. Quant. Grav.* **6** (1989) 1159.
- [12] I. Bakas, “Renormalization group flows and continual Lie algebras”, *JHEP* **0308** (2003) 013.
- [13] I. Bakas, “Ricci flows and infinite dimensional algebras”, *Fortschr. Phys.* **52** (2004) 464; “Ricci flows and their integrability in two dimensions”, *C. R. Physique* **6** (2005) 175.

- [14] M. Saveliev and A. Vershik, “Continual analogs of contragradient Lie algebras (Lie algebras with a Cartan operator and nonlinear dynamical systems)”, Commun. Math. Phys. 126 (1989) 367.
- [15] I. Bakas, “The algebraic structure of geometric flows in two dimensions”, in preparation.
- [16] M. Struwe, “A flow approach to Nirenberg’s problem”, ETH preprint (2004).
- [17] E. Glass and D. Robinson, “A nilpotent prolongation of the Robinson-Trautman equation”, J. Math. Phys. 25 (1984) 3382.
- [18] J. Finley, “The Robinson-Trautman type III prolongation structure contains K_2 ”, Commun. Math. Phys. 178 (1996) 375.
- [19] V.G. Kac, “Simple irreducible graded Lie algebras of finite growth”, Math. USSR Izv. 2 (1968) 1271; “On simplicity of certain infinite dimensional Lie algebras”, Bull. Amer. Math. Soc. 2 (1980) 311.
- [20] R. Güven and E. Yörük, “Stringy Robinson-Trautman solutions”, Phys. Rev. D54 (1996) 6413.